

## Olive Herbicide Labels & Use Restrictions

Dan Horton and Terry All, University of Georgia, Department of Entomology, Athens, GA 30602  
<http://www.ent.uga.edu/fruit.htm>

Based on labels as seen on CDMS (<http://www.cdms.net/LabelsMsds/LMDefault.aspx>) and Greenbook (<http://www.greenbook.net/>) websites (last referenced 9/24/2008). UGA provides this compilation to minimize the risk of off-label pesticide use. Always refer to the actual product label, which is the definitive guide for safe, legal pesticide use.

| Active ingredient      | WSSA group # | Trade name         | Manufacturer      | Re-Entry Interval (REI)/<br>Pre-Harvest Interval (PHI)       |
|------------------------|--------------|--------------------|-------------------|--|
| PRE EMERGENCE          |              |                    |                   |  |
| flumioxazin            | 14           | Chateau 51WDG      | Valent            | 12hrs/non-bearing only                                       |
| isoxaben               | 21           | Gallery 75DF       | Dow AgroSciences  | 12 hrs/non-bearing only                                      |
| isoxaben + trifluralin | 21 + 3       | Snapshot 2.5 TG    | Dow AgroSciences  | 12 hrs/non-bearing only                                      |
| oryzalin               | 3            | FarmSaver Oryzalin | MANA              | 24 hrs/apply late fall or early spring, orchard middles only |
|                        |              | Surflan 4AS        | United Phosphorus |  |
| oxyfluorfen            | 14           | Galigan 2E         | MANA              | 24 hrs/dormant or post-harvest                               |
|                        |              | Goal 2XL           | Dow AgroSciences  |  |
|                        |              | GoalTender 4EC     | Dow AgroSciences  |  |
| pendimethalin          | 3            | Pendulum 3.3 EC    | BASF              | 24 hrs/non-bearing only                                      |
| simazine               | 5            | Princep 4L         | Syngenta          | 12 hrs/apply late fall or mid-winter                         |
|                        |              | Simazine 4L        | Agrisolutions     |  |
| POST EMERGENCE         |              |                    |                   |  |
| carfentrazone          | 14           | Aim 2EC            | FMC               | 12 hrs/3 days  |
| clethodim              | 1            | Arrow 2 EC         | Makhteshim-Agan   | 24 hrs/non-bearing only                                      |
|                        |              | Clethodim 2E       | Albaugh           |  |
|                        |              | Envoy Herbicide    | Valent            |  |
|                        |              | Select 2EC         | Valent            |  |
|                        |              | Select Max         | Valent            |  |
| fluazifop              | 1            | Fusilade DX        | Syngenta          | 12 hrs/non-bearing only                                      |
| glyphosate             | 9            | Cinco              | Loveland          | 12 hrs/directed sprays only, 17 days                         |
|                        |              | Roundup WeatherMax | Monsanto          | 4 hrs/directed sprays only, NLT 90 days post-bloom           |
| paraquat dichloride    | 22           | Firestorm          | Chemtura          | 12 to 24 hrs/directed sprays only, 13 days                   |
|                        |              | Gramoxone Inteon   | Syngenta          |  |
| sethoxydim             | 1            | Poast              | BASF              | 12 hrs/non-bearing only                                      |
| clove oil              |              | Matran EC Organic  | EcoSMART          | when dry/when dry  |
| pelargonic acid        |              | Scythe             | Dow AgroSciences  | 12 hrs/24 hrs  |
| benefin + oryzalin     | 3 + 3        | XL 2G              | Helena Chemical   | 24 hrs/non-bearing only                                      |